

Contamination of short GRBs by giant magnetar flares: Significance of downward revision in distance to SGR 1806–20

Paul A. Crowther^a, Joanne L. Bibby^a, James P. Furness^a, J. Simon Clark^b

^aDept of Physics & Astronomy, University of Sheffield, Sheffield, S3 7RH, UK

^bDept of Physics & Astronomy, The Open University, Milton Keynes, MK7 6AA, UK

We highlight how the downward revision in the distance to the star cluster associated with SGR 1806–20 by Bibby et al. (2008) reconciles the apparent low contamination of BATSE short GRBs by intense flares from extragalactic magnetars without recourse to modifying the frequency of one such flare per 30 years per Milky Way galaxy. We also discuss the variety in progenitor initial masses of magnetars based upon cluster ages, ranging from $\sim 50M_{\odot}$ for SGR 1806–20 and AXP CXOU J164710.2-455216 in Westerlund 1 to $\sim 17M_{\odot}$ for SGR 1900+14 according to Davies et al. (2009) and presumably also 1E 1841–045 if it originated from one of the massive RSG clusters #2 or #3.

1. INTRODUCTION

It is well known that the initial, intense γ -ray spike from giant flares of extragalactic magnetars – locally classified as either Soft Gamma Repeaters (SGRs) or Anomalous X-ray Pulsars (AXPs) at kpc distances – could be mistaken for cosmological short Gamma Ray Bursts (GRBs) (Hurley et al. 2005). Typical peak outputs of magnetar flares lie in the range $10^{41} \text{ erg s}^{-1}$ (SGR 1806–20 in Jan 1979) to $10^{45} \text{ erg s}^{-1}$ (SGR 0525–66 in Mar 1979) would not be detected in galaxies lying at Mpc distances, but the peak luminosity of the intense flare from SGR 1806–20 in Dec 2004 was $2 \times 10^{47} d_{15} \text{ erg s}^{-1}$ where the usually adopted distance of SGR 1806–20 is $d_{15} = 15 \text{ kpc}$. BATSE, aboard the Compton Gamma-Ray Observatory, would have detected the initial $\sim 0.1 \text{ s}$ spike from SGR 1806–20 had it originated from a galaxy $30\text{--}50 d_{15} \text{ Mpc}$ away.

For an adopted rate of one such intense flare, per 30 yr, per Milky Way galaxy, a total of 200–300 $d_{15}^3 (\tau/30\text{yr})^{-1}$ flares would have been detected by BATSE over its 9.5 year lifetime, representing 40–60% $d_{15}^3 (\tau/30\text{yr})^{-1}$ of all short GRBs detected by BATSE (Hurley et al. 2005; Lazzati et al. 2005). Consequently, one would expect a strong correlation between the position of BATSE

short bursts and nearby star-forming galaxies. However, Tanvir et al. (2005) found that only 10% of short GRBs originated from within 30 Mpc, while Lazzati et al. (2005) concluded that the contribution of giant magnetar flares to short GRB statistics was no greater than $\sim 4\%$. Only a few candidate extragalactic giant flares have been proposed, such as GRB051103 in M81 (Ofek et al. 2006), although such claims remain inconclusive (Hurley et al. 2010). Apparently, the frequency of intense magnetar flares is much lower than one per 30 years, which remains a very credible possibility, unless the widely adopted 15 kpc distance to SGR 1806–20 is incorrect.

Fortunately, SGR 1806–20 lies towards a young star cluster, whose near-IR appearance is shown in Fig. 1. Photometric (colour-magnitude diagrams) and spectroscopic (stellar subtypes) tools allow ages and main-sequence turn-off masses to be derived, albeit somewhat dependent upon isochrones from evolutionary models of massive stars. Walborn (2009) discusses the characteristics of the youngest star clusters, for which O stars represent the visually brightest stars up to ages of a few Myr (e.g. NGC 3603) with AF supergiants dominating for ages closer to 4–10 Myr (e.g. Westerlund 1). Thereafter, the visual and near-IR appearance of star clusters is distin-




Figure 1. Location of magnetars SGR 1806–20 and AXP CXOU J164710.2–455216 within visibly obscured young massive clusters: *Left* Westerlund 1 (NTT/SOFI, 200×200 arcsec, equivalent to 5×5 pc at $d=5$ kpc); *right* Cl 1806–20 (Gemini-N/NIRI, 34×34 arcsec, equivalent to 1.5×1.5 pc at $d=8.7$ kpc).

guished by red supergiants (e.g. RSG cluster #1: Figer et al. 2006; RSG cluster #2: Davies et al. 2007).

2. DISTANCE TO SGR 1806–20

The usual 15 kpc distance towards SGR 1806–20 relies upon kinematics (Corbel et al. 1997; Eikenberry et al. 2004; Figer et al. 2004), but if we assume that it is physically associated with a visibly obscured ($A_K = 3$ mag), massive star cluster (Eikenberry et al. 2004; Figer et al. 2005), we may combine their physical properties with evolutionary models to obtain the age and (spectroscopic parallax) distance to the cluster. The likelihood of chance alignment between the cluster and magnetar is low. Indeed, Munro et al. (2006) obtained a confidence of $>99.97\%$ for the association of magnetar CXOU J164710.2–455217 (AXP) with the Westerlund 1 cluster (see Fig. 1). Both clusters host rare Luminous Blue Variables and Wolf-Rayet stars in addition to OB stars, although Westerlund 1 is an order of magnitude more massive ($5\text{--}10 \times 10^4 M_\odot$; Clark et al. 2005; Mengel & Tacconi-Garman 2007; Brandner et al. 2008), so hosts many more evolved massive stars – 24 WR stars (Crowther et al. 2006b) versus 4 WR stars in Cl 1806–20 (Eikenberry et al. 2004; Figer et al. 2005).

Bibby et al. (2008) presented Gemini-S/GNIRS near-IR spectroscopy of OB supergiants and non-dusty Wolf-Rayet stars in SGR 1806–20 from which a cluster distance of $8.7^{+1.8}_{-1.5}$ kpc was obtained using a calibration of spectral type versus absolute magnitude. Reliable OB spectral types were obtained in two cases from comparison with the near-IR atlas of Hanson et al. (2005). Dust producing WR stars are highly unreliable as absolute magnitude calibrators since their near-IR appearance is dictated by the properties of their dust rather than underlying stellar continua (Crowther et al. 2006b). Main sequence OB stars would provide superior absolute magnitude calibrators with respect to Wolf-Rayet stars and OB supergiants, but these would require long near-IR integrations for reliable stellar classification even for present 8–10m telescopes.

Alternatively, the simultaneous presence of




Figure 2. Meynet et al. (1994) isochrone fits (shown in black) to properties of B supergiants in SGR 1806–20 (shown in red from Bibby et al. 2008) for ages of 2.8 Myr (top), 4 Myr (middle) and 5 Myr (bottom), from which distances of 15, 8.7 and 6.3 kpc are obtained. For the conventional 15 kpc distance to SGR 1806–20, the cluster OB stars would exhibit hypergiant spectral morphologies, while WC stars are not predicted to exist at such young ages.




Figure 3. Comparison between various magnetar and cluster distances to 1806-20, uniformly adapted to a Galactic Centre distance of 8 kpc (Reid 1993). Updated from Bibby et al. (2008).

WN and WC-type Wolf-Rayet stars in SGR 1806-20 suggest an age of 4 ± 1 Myr, from which an independent distance measurement was obtained using isochrones for OB stars (Lejeune & Schaerer 2001) based on the evolutionary models of Meynet et al. (1994) and the B supergiant temperature calibration of Crowther et al. (2006a). For ages of 2.8, 4 and 5 Myr, both cluster distances and magnetar masses were obtained, as presented in Fig 2. Let us consider the properties of B supergiants and WC stars for each of the three cases in turn,

- **15 kpc** For this distance, the cluster B-type stars studied by Bibby et al. (2008) would be exceptionally luminous, with four of the five cases exceeding $\log(L/L_{\odot}) = 6$. Instances of such luminous B supergiants are known, such as ζ^1 Sco (Crowther et al. 2006a). However, such stars display hypergiant spectral morphologies (B1.5 Ia⁺ for ζ^1 Sco), whereas the B stars in CI 1806-20 appear to be morphologically normal Ia, Iab or Ib supergiants. In addition, solar metallicity evolutionary models for 120

$M_{\odot}$ stars allowing for rotational mixing and contemporary mass-loss prescriptions indicate that the WC phase is not predicted until after 3.5 Myr for a model initially rotating at 40% of critical (maximum) velocity (Hirschi, priv. comm.). For the 120 $M_{\odot}$ non-rotating case, the WC phase would be anticipated to *commence* after 2.8 Myr (Meynet & Maeder 2003).

- **8.7 kpc** For this distance, the cluster B-type stars would be possess luminosities in the range $\log(L/L_{\odot}) = 5.4 - 5.7$, close to the $\log(L/L_{\odot}) = 5.5$ average of 25 O9.5-B3 Ia supergiants studied by Crowther et al. (2006a), the majority of which are morphologically normal. From Meynet & Maeder (2003), the hydrogen-burning phase of a non-rotating 50 $M_{\odot}$ star is predicted to last for 4.0 Myr, followed by short WN and WC phases, each of ~ 0.1 Myr. In this case WC stars would be expected after ~ 4.1 Myr. Models initially rotating at 300 km s^{-1} suggest longer main-sequence and WR phases, corresponding to the onset of the WC phase

after 5 Myr.

- **6.3 kpc** For this distance, the cluster B-type stars would be possess luminosities in the range $\log(L/L_\odot) = 5.1 - 5.5$, somewhat below the average of the Ia supergiants studied by Crowther et al. (2006a), and more representative of Iab or Ib supergiants (Searle et al. 2008). Our GNIRS near-IR spectroscopic datasets does not allow us to discriminate between Ia and Ib supergiants. Again, turning to the WR population, from Meynet & Maeder (2003), a non-rotating $40 M_\odot$ solar metallicity model would not be expected to remain as a WN star prior to core-collapse, whereas a model initially rotating at 300 km s^{-1} is predicted to advance to the WC phase only after ~ 5.7 Myr.

Overall, a spectroscopic distance of 7–10 kpc is favoured from the simultaneous presence of normal B supergiants, plus WN and WC stars in Cl 1806–20. This cluster distance is compared to previous work in Fig. 3, and is consistent with both the Cameron et al. (2005) magnetar distance and McClure-Griffiths & Gaensler (2005) reanalysis thereof.

3. CONTRIBUTION OF EXTRA-GALACTIC GIANT FLARES TO BATSE SHORT BURSTS

Overall, we find a reduced cluster distance of $d_{15} = 0.58 \pm 0.1$, and since the contribution of giant flares to BATSE short GRB statistics scales with d_{15}^3 we find that giant flares contribute $8^{+5}_{-4}\%$ of all BATSE short bursts, for the *canonical* rate of one giant flare per 30 yr per Milky Way galaxy. Of course, the frequency of once per 30 years is merely a crude upper limit, since it originates from the fact that solely one giant flare (from SGR 1806–20) has been detected from a Milky Way source since the advent of gamma-ray telescopes. Nevertheless, our result largely resolves the previous (low) rate problem, namely (i) $\sim 4\%$ from the scarcity of BATSE sources with spectral properties similar to giant flares (Lazzati et al. 2005); (ii) $\sim 10\%$ from giant flares within 30 Mpc

(Tanvir et al. 2005); (iii) $< 15\%$ from the lack of local host galaxies for several well-localized short GRBs (Nakar et al. 2006); (iv) a few percent contamination from the absence of giant flares from galaxies in the Virgo cluster (Popov & Stern 2006).

4. MAGNETAR MASSES FROM ASSOCIATED CLUSTERS

As discussed above, associated clusters allow progenitor masses of magnetars to be estimated, since magnetars are believed to be young objects. For SGR 1806–20, Bibby et al. (2008) infer a progenitor mass of $48^{+20}_{-8} M_\odot$, adding to evidence linking some magnetars to very massive stars. Identical conclusions were reached by Munro et al. (2006) for the AXP CXOU J164710.2-455216 in the star cluster Westerlund 1 (Clark et al. 2005) whose age and stellar content is remarkably similar to Cl 1806–20. Other observations have also been used to link magnetars to high mass progenitors. For example, Gaensler et al. (2005) have interpreted a hydrogen shell centred upon the AXP 1E 1048.1–5937 as a wind blown bubble from its 30–40 $M_\odot$ progenitor.

However, lower progenitor mass magnetars are also known. SGR 1900+14 possesses a pair of M5 supergiants (Vrba et al. 1996; 2000). Davies et al. (2009) have recently presented deep K-band imaging of these red supergiants, in which faint members of the associated cluster Cl 1900+14 can be identified. Davies et al. (2009) have obtained a kinematic distance of 12.5 kpc to these M supergiants, and an evolutionary age of 14 ± 1 Myr, based upon contemporary solar metallicity evolutionary models of Meynet & Maeder (2000) plus bolometric corrections of Levesque et al. (2005). If one assumes that the progenitor of SGR 1900+14 originated from this cluster, an initial mass of $17 \pm 2 M_\odot$ can be deduced. Similar results would be obtained for the AXP 1E 1841–045 (within Kes 73) *if* it originated from either RSG cluster #2 (Davies et al. 2007) or RSG cluster #3 (Clark et al. 2009). These are located at a similar distance to 1E 1841–045, within a few degrees of its sight-line, although the likelihood of physical association is naturally much weaker

Table 1
Progenitor masses of magnetars based on associated star clusters.

SGR/AXP	Cluster	d kpc	Age Myr	Mass $M_{\odot}$	Reference
1806-20	Cl 1806-20	$8.7^{+1.8}_{-1.5}$	4 ± 1	48^{+20}_{-8}	Bibby et al. (2008)
CXOU J164710.2-455216	Westerlund 1	5	~ 4.5	~ 55	Clark et al. (2008)
1900+14	Cl 1900+14	12.5	14 ± 1	17 ± 2	Davies et al. (2009)
1E 1841-045	RSGC #2 or #3	6	16 ± 4	~ 15	Clark et al. (2009)

than for SGR 1806-20, CXOU J164710.2-455216 and SGR 1900+14.

Table 1 provides a summary of magnetar progenitor masses inferred from potentially associated star clusters. In the Milky Way, it is apparent that magnetars can be produced by both $\sim 15M_{\odot}$ and $\sim 50M_{\odot}$ stars. The former presumably originated in Type II-P SN following the red supergiant (RSG) phase, while the latter are likely to have originated in a Type Ibc SN following the WR phase. From comparison with Heger et al. (2003), certainly SGR 1900+14 and possibly SGR 1806-20 would be expected to produce neutron star remnants at Solar metallicity. For metal poor populations the latter channel is unlikely to be available since reduced mass-loss rates during the main sequence and post-main sequence phases would likely lead to the formation of a black hole for such high initial mass stars. No direct evidence for supernova remnants is known in Cl 1806-20, Westerlund 1 or Cl 1900+14. Presumably this is because the combined stellar winds from other cluster members will blow a sufficiently large cavity that a supernova remnant could expand into without showing any observable emission.

Finally, it should be borne in mind that cluster ages and magnetar progenitor masses rely upon the reliability of evolutionary models. Values quoted herein are based upon a set of assumptions which have subsequently been improved upon through mass-loss rate prescriptions and allowance for rotational mixing (Meynet & Maeder 2000). Close binary evolution also further complicates inferred main-sequence turn-off masses. Therefore, one should not necessarily treat specific progenitor masses or cluster ages as robust, although the clear differences between the stel-

lar content of clusters 1806-20 and 1900+14 undoubtedly demonstrate distinct channels leading to the production of magnetars.

REFERENCES

1. Bibby, J.L., Crowther, P.A., Furness, J.P., Clark, J.S. A downward revision to the distance of the 1806-20 cluster and associated magnetar from Gemini Near-Infrared Spectroscopy, *Mon. Not. R. Astron. Soc.*, 386, L23-L27, 2008.
2. Brandner, W., Clark, J.S., Stolte, A. et al. Intermediate to low-mass stellar content of Westerlund 1, *Astron. & Astrophys.*, 478, 137-149, 2008.
3. Cameron P.B., Chandra, P., Ray, A. et al. Detection of a radio counterpart to the 27 December 2004 giant flare from SGR 1806 - 20, *Nat.* 434, 1112-1115, 2005.
4. Clark J.S., Negueruela, I., Crowther, P.A., Goodwin, S.P. On the massive stellar population of the super star cluster Westerlund 1, *Astron. & Astrophys.*, 434, 949-969, 2005.
5. Clark J.S., Munro, M.P., Negueruela, I. et al. Unveiling the X-ray point source population of the Young Massive Cluster Westerlund 1, *Astron. & Astrophys.*, 477, 147-163, 2008.
6. Clark J.S., Negueruela, I., Davies, B. et al. A third red supergiant rich cluster in the Scutum-Crux arm, *Astron. & Astrophys.* 498, 109-114, 2009.
7. Corbel S., Wallyn, O., Dame, T.M. et al. The Distance of the Soft Gamma Repeater SGR 1806-20, *Astrophys. J.*, 478, 624-630, 1997.
8. Crowther P.A., Lennon, D.J., Walborn, N.R., Physical parameters and wind properties of galactic early B supergiants, *Astron. & As-*

- trophys.*, 446, 279–293, 2006a.
9. Crowther P.A., Hadfield, L.J., Clark, J.S. et al. A census of the Wolf-Rayet content in Westerlund 1 from near-infrared imaging and spectroscopy, *Mon. Not. Roy. Astron. Soc.*, 372, 1407–1424, 2006b.
 10. Davies B., Figer, D.F., Kudritzki, R.-P. et al. A Massive Cluster of Red Supergiants at the Base of the Scutum-Crux Arm, *Astrophys. J.*, 671, 781–801, 2007.
 11. Davies B., Figer, D.F., Kudritzki, R.-P. et al. The progenitor mass of the magnetar SGR 1900+14. *Astrophys. J.*, 707, 844–851, 2009.
 12. Eikenberry S.S., Matthews, K., LaVine, J.L. et al. Infrared Observations of the Candidate LBV 1806-20 and Nearby Cluster Stars, *Astrophys. J.*, 616, 506–518, 2004.
 13. Figer D.F., Najarro, F., Kudritzki, R.-P. The Double-lined Spectrum of LBV 1806-20, *Astrophys. J.*, 610, L109–L112, 2004.
 14. Figer D.F. Najarro, F., Geballe, T.R. et al. Massive Stars in the SGR 1806-20 Cluster, *Astrophys. J.*, 622, L49–L52, 2005. bib-item Figer06 Figer D.F., MacKenty, J.W., Roberto, M. et al. Discovery of an Extraordinarily Massive Cluster of Red Supergiants, *Astrophys. J.*, 643, 1166–1179, 2006.
 15. Gaensler, B.M., McClure-Griffiths, N.M., Oey, M.E. et al. A stellar wind bubble coincident with the Anomalous X-ray pulsar 1E 1048.1–5937: Are magnetars formed from massive progenitors? *Astrophys. J.*, 620, L95–L99, 2005.
 16. Hanson, M.M., Kudritzki, R.-P., Kenworthy, M.A. et al. A Medium Resolution Near-Infrared Spectral Atlas of O and Early-B Stars, *Astrophys. J. Suppl.*, 161, 154–170, 2005.
 17. Heger A., Fryer, C.L., Woosley, S.E. et al. How Massive Single Stars End Their Life, *Astrophys. J.*, 591, 288–300, 2003.
 18. Hurley, K., Boggs, S.E., Smith, D.M. et al. An exceptionally bright flare from SGR 1806-20 and the origins of short-duration γ -ray bursts, *Nat.* 434, 1098–1103, 2005.
 19. Hurley, K., Rowlinson, A., Bellm, E. et al. A new analysis of the short-duration, hard-spectrum GRB 051103, a possible extragalactic SGR giant flare, *Mon. Nat. Roy. Astron. Soc.* submitted (arXiv:0907.2462)
 20. Lazzati D., Ghirlanda, G., Ghisellini, G. Soft gamma-ray repeater giant flares in the BATSE short gamma-ray burst catalogue: constraints from spectroscopy, *Mon. Not. Roy. Astron. Soc.*, 362, L8–L12, 2005.
 21. Lejeune, T., Schaerer, D., Database of Geneva stellar evolution tracks and isochrones for (UBV)_J(RI)CJHKLL'M, HST-WFPC2, Geneva and Washington photometric systems, *Astron. & Astrophys.*, 366, 538–546, 2001.
 22. Levesque, E.M., Massey, P.M., Olsen, K.A.G. et al. The Effective Temperature Scale of Galactic Red Supergiants: Cool, but Not As Cool As We Thought, *Astrophys. J.*, 628, 973–985, 2005.
 23. McClure-Griffiths, N.M., Gaensler, B.M. Constraints on the Distance to SGR 1806-20 from H I Absorption, *Astrophys. J.* 630, L161–L163, 2005.
 24. Mengel, S., Tacconi-Garman, L.E. Medium resolution 2.3 μ m spectroscopy of the massive Galactic open cluster Westerlund 1, *Astron. & Astrophys.* 466, 151–155, 2007.
 25. Meynet G., Maeder A., Stellar evolution with rotation. V. Changes in all the outputs of massive star models, *Astron. & Astrophys.* 361, 101–120, 2000.
 26. Meynet G., Maeder A., Stellar evolution with rotation. X. Wolf-Rayet star populations at solar metallicity, *Astron. & Astrophys.* 404, 975–990, 2003.
 27. Muno, M.P., Clark, J.S., Crowther, P.A. et al. A Neutron Star with a Massive Progenitor in Westerlund 1, *Astrophys. J.*, 636, L41–L44, 2006.
 28. Nakar, E., Gal-Yam, A., Piran, T., Fox, D.B. The Distances of Short-Hard Gamma-Ray Bursts and the Soft Gamma-Ray Repeater Connection, *Astrophys. J.*, 640, 849–83, 2006.
 29. Ofek, E.O. Kulkarni, S.R., Nakar, E. et al. The Short-Hard GRB 051103: Observations and Implications for Its Nature, *Astrophys. J.*, 652, 507–511, 2006.
 30. Popov, S.B., Stern, B.E., Soft gamma re-

- peaters outside the Local Group, *Mon. Nat. Roy. Astron. Soc.*, 365, 885-890, 2006.
31. Reid, M.J., The distance to the center of the Galaxy, *Ann. Rev. Astron. & Astrophys.*, 31, 345-372, 1993.
 32. Searle, S.C., Prinja, R.K., Massa, D., Ryans, R. Quantitative studies of the optical and UV spectra of Galactic early B supergiants I. Fundamental parameters, *Astron. Astrophys.* 481, 777-797, 2008.
 33. Tanvir N.R., Chapman, R., Levan, A.J., Pridey, R.S. An origin in the local Universe for some short γ -ray bursts, *Nat.*, 438, 991-993, 2005.
 34. Vrba, F.J., Luginbuhl, C.B., Hurley, K.C. et al. The Double Infrared Source toward the Soft Gamma-Ray Repeater SGR 1900+14. *Astrophys. J.*, 648, 225-230, 1996.
 35. Vrba, F.J., Henden, A.A., Luginbuhl, C.B. et al. The Discovery of an Embedded Cluster of High-Mass Stars near SGR 1900+14, *Astrophys. J.*, 533, L17-L20, 2000.
 36. Walborn, N.R., Age Paradigms for Massive Young Clusters, in Proc: *Hot and Cool: Bridging Gaps in Massive Star Evolution*, ASP Conf. Ser (eds. C. Leitherer et al.), in press, 2009.